arXiv:1511.00948v2 [math.NT] 4 Nov 2015

# INTEGER SETS WITH IDENTICAL REPRESENTATION FUNCTIONS

YONG-GAO CHEN AND VSEVOLOD F. LEV

Abstract. We present a versatile construction allowing one to obtain pairs of integer sets with infinite symmetric difference, infinite intersection, and identical representation functions.

Let $\mathbb{N}_0$ denote the set of all non-negative integers. To every subset $A \subseteq \mathbb{N}_0$ corresponds its representation function $R_A$ defined by

$$R_A(n) := |\{(a', a'') \in A \times A \colon n = a' + a'',\ a' < a''\}|;$$

that is, $R_A(n)$ is the number of unordered representations of the integer $n$ as a sum of two distinct elements of $A$.

Answering a question of Sárközy, Dombi [D02] constructed sets $A, B \subseteq \mathbb{N}_0$ with infinite symmetric difference such that $R_A = R_B$. The result of Dombi was further extended and developed in [CW03] (where a different representation function was considered) and [L04] (a simple common proof of the results from [D02] and [CW03] using generating functions); other related results can be found in [C11, CT09, Q15, T08].

The two sets constructed by Dombi actually *partition* the ground set $\mathbb{N}_0$, which makes one wonder whether one can find $A, B \subseteq \mathbb{N}_0$ with $R_A = R_B$ so that not only the symmetric difference of $A$ and $B$, but also their intersection is infinite. Tang and Yu [TY12] proved that if $A \cup B = \mathbb{N}_0$ and $R_A(n) = R_B(n)$ for all sufficiently large integer $n$, then at least one cannot have $A \cap B = 4\mathbb{N}_0$ (here and below $k\mathbb{N}_0$ denotes the dilate of the set $\mathbb{N}_0$ by the factor $k$). They further conjectured that, indeed, under the same assumptions, the intersection $A \cap B$ cannot be an infinite arithmetic progression, unless $A = B = \mathbb{N}_0$. The main goal of this note is to resolve the conjecture of Tang and Yu in the negative by constructing an infinite family of pairs of sets $A, B \subseteq \mathbb{N}_0$ with $R_A = R_B$ such that $A \cup B = \mathbb{N}_0$, while $A \cap B$ is an infinite arithmetic progression properly contained in $\mathbb{N}_0$. Our method also allows one to easily construct sets $A, B \subseteq \mathbb{N}_0$ with $R_A = R_B$ such that both their symmetric difference and intersection are infinite, while their union is arbitrarily sparse and the intersection is *not* an arithmetic progression.

2010 *Mathematics Subject Classification.* 11B34.
*Key words and phrases.* Representation function.

For sets $A, B \subseteq \mathbb{N}_0$ and integer $m$ let $A - B := \{a - b \colon (a, b) \in A \times B\}$ and $m + A := \{m + a \colon a \in A\}$.

The following basic lemma is in the heart of our construction.

**Lemma 1.** *Suppose that $A_0, B_0 \subseteq \mathbb{N}_0$ satisfy $R_{A_0} = R_{B_0}$, and that $m$ is a non-negative integer with $m \notin (A_0 - B_0) \cup (B_0 - A_0)$. Then, letting*

$$A_1 := A_0 \cup (m + B_0) \text{ and } B_1 := B_0 \cup (m + A_0),$$

*we have $R_{A_1} = R_{B_1}$ and furthermore*

i) $A_1 \cup B_1 = (A_0 \cup B_0) \cup (m + A_0 \cup B_0)$;
ii) $A_1 \cap B_1 \supseteq (A_0 \cap B_0) \cup (m + A_0 \cap B_0)$, *the union being disjoint.*

*Moreover, if $m \notin (A_0 - A_0) \cup (B_0 - B_0)$, then also in i) the union is disjoint, and in ii) the inclusion is in fact an equality.*

*In particular, if $A_0 \cup B_0 = [0, m-1]$, then $A_1 \cup B_1 = [0, 2m-1]$, and if $A_0$ and $B_0$ indeed* partition *the interval $[0, m-1]$, then $A_1$ and $B_1$ partition the interval $[0, 2m-1]$.*

*Proof.* Since the assumption $m \notin A_0 - B_0$ ensures that $A_0$ is disjoint from $m + B_0$, for any integer $n$ we have

$$R_{A_1}(n) = R_{A_0}(n) + R_{B_0}(n - 2m) + |\{(a_0, b_0) \in A_0 \times B_0 \colon a_0 + b_0 = n - m\}|.$$

Similarly,

$$R_{B_1}(n) = R_{B_0}(n) + R_{A_0}(n - 2m) + |\{(a_0, b_0) \in A_0 \times B_0 \colon a_0 + b_0 = n - m\}|,$$

and in view of $R_{A_0} = R_{B_0}$, this gives $R_{A_1} = R_{B_1}$. The remaining assertions are straightforward to verify. □

Given subsets $A_0, B_0 \subseteq \mathbb{N}_0$ and a sequence $(m_i)_{i \in \mathbb{N}_0}$ with $m_i \in \mathbb{N}_0$ for each $i \in \mathbb{N}_0$, define subsequently

$$A_i := A_{i-1} \cup (m_{i-1} + B_{i-1}) \text{ and } B_i := B_{i-1} \cup (m_{i-1} + A_{i-1}), \quad i = 1, 2, \dots \tag{1}$$

and let

$$A := \cup_{i \in \mathbb{N}_0} A_i, \ B := \cup_{i \in \mathbb{N}_0} B_i. \tag{2}$$

As an immediate corollary of Lemma 1, if $R_{A_0} = R_{B_0}$ and $m_i \notin (A_i - B_i) \cup (B_i - A_i)$ for each $i \in \mathbb{N}_0$, then $R_A = R_B$.

The special case $A_0 = \{0\}$, $B_0 = \{1\}$, $m_i = 2^{i+1}$ yields the partition of Dombi (which, we remark, was originally expressed in completely different terms). Below we analyze yet another special case obtained by fixing arbitrarily an integer $l \geq 1$ and

choosing $A_0 := \{0\}$, $B_0 := \{1\}$, and

$$m_i := \begin{cases} 2^{i+1}, & 0 \le i \le 2l-2, \\ 2^{2l}-1, & i = 2l-1, \\ 2^{i+1}-2^{i-2l}, & i \ge 2l. \end{cases} \tag{3}$$

We notice that $R_{A_0} = R_{B_0}$ in a trivial way (both functions are identically equal to 0), and that $A_0$ and $B_0$ partition the interval $[0, m_0 - 1]$. Applying Lemma 1 inductively $2l-2$ times, we conclude that in fact for each $i \le 2l-2$, the sets $A_i$ and $B_i$ partition the interval $[0, 2m_{i-1} - 1] = [0, m_i - 1]$, and consequently $m_i \notin (A_i - B_i) \cup (B_i - A_i)$ and $m_i \notin (A_i - A_i) \cup (B_i - B_i)$. In particular, $A_{2l-2}$ and $B_{2l-2}$ partition $[0, m_{2l-2} - 1]$, and therefore $A_{2l-1}$ and $B_{2l-1}$ partition $[0, 2m_{2l-2} - 1] = [0, m_{2l-1}]$. In addition, it is easily seen that $A_{2l-1}$ contains both 0 and $m_{2l-1}$, whence $m_{2l-1} \in A_{2l-1} - A_{2l-1}$, but $m_{2l-1} \notin B_{2l-1} - B_{2l-1}$ and $m_{2l-1} \notin (A_{2l-1} - B_{2l-1}) \cup (B_{2l-1} - A_{2l-1})$. From Lemma 1 i) it follows now that $A_{2l} \cup B_{2l} = [0, 2m_{2l-1}] = [0, m_{2l} - 1]$, while

$$A_{2l} \cap B_{2l} = \Big(A_{2l-1} \cap (m_{2l-1} + A_{2l-1})\Big) \cup \Big(B_{2l-1} \cap (m_{2l-1} + B_{2l-1})\Big) = \{m_{2l-1}\}.$$

Applying again Lemma 1 we then conclude that for each $i \ge 2l$,

$$A_i \cup B_i = [0, m_i - 1]$$

(implying $m_i \notin (A_i - B_i) \cup (B_i - A_i)$) and

$$A_i \cap B_i = m_{2l-1} + \{0, m_{2l}, 2m_{2l}, \dots, (2^{i-2l} - 1)m_{2l}\}.$$

As a result, with $A$ and $B$ defined by (2), we have $A \cup B = \mathbb{N}_0$ while the intersection of $A$ and $B$ is the infinite arithmetic progression $m_{2l-1} + m_{2l}\mathbb{N}_0$. Moreover, the condition $m_i \notin (A_i - B_i) \cup (B_i - A_i)$, which we have verified above to hold for each $i \ge 0$, results in $R_A = R_B$.

We thus have proved

**Theorem 1.** *Let $l$ be a positive integer, and suppose that the sets $A, B \subseteq \mathbb{N}_0$ are obtained as in (1)–(2) starting from $A_0 = \{0\}$ and $B_0 = \{1\}$, with $(m_i)$ defined by (3). Then $R_A = R_B$, while $A \cup B = \mathbb{N}_0$ and $A \cap B = (2^{2l} - 1) + (2^{2l+1} - 1)\mathbb{N}_0$.*

We notice that for any fixed integers $r \ge 2^{2l} - 1$ and $m \ge 2^{2l+1} - 1$, having (3) appropriately modified (namely, setting $m_i = 2^{i-2l}m$ for $i \ge 2l$) and translating $A$ and $B$, one can replace the progression $(2^{2l} - 1) + (2^{2l+1} - 1)\mathbb{N}_0$ in the statement of Theorem 1 with the progression $r + m\mathbb{N}_0$; however, the relation $A \cup B = \mathbb{N}_0$ will *not* hold true any longer unless $r = 2^{2l} - 1$ and $m = 2^{2l+1} - 1$. This suggests the following question.

*Problem* 1. Given that $R_A = R_B$, $A \cup B = \mathbb{N}_0$, and $A \cap B = r + m\mathbb{N}_0$ with integer $r \ge 0$ and $m \ge 2$, must there exist an integer $l \ge 1$ such that $r = 2^{2l} - 1$, $m = 2^{2l+1} - 1$, and $A, B$ are as in Theorem 1?

The finite version of this question is as follows.

*Problem* 2. Given that $R_A = R_B$, $A \cup B = [0, m-1]$, and $A \cap B = \{r\}$ with integer $r \geq 0$ and $m \geq 2$, must there exist an integer $l \geq 1$ such that $r = 2^{2l}-1$, $m = 2^{2l+1}-1$, $A = A_{2l}$, and $B = B_{2l}$, with $A_{2l}$ nd $B_{2l}$ as in the proof of Theorem 1?

We conclude our note with yet another natural problem.

*Problem* 3. Do there exist sets $A, B \subseteq \mathbb{N}_0$ with the infinite symmetric difference and with $R_A = R_B$ which *cannot* be obtained by a repeated application of Lemma 1?

**Acknowledgments.** Early stages of our work depended heavily on extensive computer programming that was kindly carried out for us by Talmon Silver; we are indebted to him for this contribution.

The first author is supported by the National Natural Science Foundation of China, Grant No. 11371195, and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

## References

[C11] Y.G. Chen, On the values of representation functions, *Sci. China Math.* **54** (2011), 1317–1331.

[CT09] Y.G. Chen and M. Tang, Partitions of natural numbers with the same representation functions, *J. Number Theory* **129** (2009), 2689–2695.

[CW03] Y.G. Chen and B. Wang, On additive properties of two special sequences, *Acta Arith.* **110** (2003), 299–303.

[D02] G. Dombi, Additive properties of certain sets, *Acta Arith.* **103** (2002), 137–146.

[L04] V.F. Lev, Reconstructing integer sets from their representation functions, *Electron. J. Combin.* **11** (2004), paper #R78.

[Q15] Z. Qu, On the nonvanishing of representation functions of some special sequences, *Discrete Math.* **338** (2015), 571–575.

[S04] C. Sándor, Partitions of natural numbers and their representation functions, *Integers* **4** (2004), paper #A18.

[T08] M. Tang, Partitions of the set of natural numbers and their representation functions, *Discrete Math.* **308** (2008), 2614–2616.

[TY12] M. Tang and W. Yu, A note on partitions of natural numbers and their representation functions, *Integers* **12** (2012), paper #A53.

School of Mathematical Sciences and Institute of Mathematics, Nanjing Normal University, Nanjing 210023, China

*E-mail address*: ygchen@njnu.edu.cn

Department of Mathematics, The University of Haifa at Oranim, Tivon 36006, Israel

*E-mail address*: seva@math.haifa.ac.il